

Tiling Lattices with Sublattices, I

David Feldman, University of New Hampshire
 James Propp, University of Massachusetts Lowell
 Sinai Robins, Nanyang Technological University

ABSTRACT: Call a coset C of a subgroup of $\mathbf{Z}^d$ a *Cartesian coset* if C equals the Cartesian product of d arithmetic progressions. Generalizing Mirsky-Newman, we show that a non-trivial disjoint family of Cartesian cosets with union $\mathbf{Z}^d$ always contains two cosets that differ only by translation. Where Mirsky-Newman's proof (for $d = 1$) uses complex analysis, we employ Fourier techniques. Relaxing the Cartesian requirement, for $d > 2$ we provide examples where $\mathbf{Z}^d$ occurs as the disjoint union of four cosets of distinct subgroups (with one not Cartesian). Whether one can do the same for $d = 2$ remains open.

In 1950, Paul Erdős conjectured that if a system of k arithmetic progressions, with distinct differences n_i , covers the natural numbers $\mathbf{N}$ (or the integers $\mathbf{Z}$), then

$$\sum_{i=1}^k \frac{1}{n_i} > 1 .$$

Erdős credits Leon Mirsky and Donald Newman with the first proof (and also cites Harold Davenport and Richard Rado for independently finding the same proof just a bit later). Mirsky and Newman did not publish at the time, but Erdős credits them in a 1952 paper. For Newman's own exposition, see his analytic number theory monograph. For more history, see Soifer. The Mirsky-Newman proof blends formal power series and complex analysis, and now serves as a standard example showing how such interactions solve combinatorial problems.

Nowadays one usually hears the Mirsky-Newman stated in the contrapositive: a non-trivial finite disjoint family of arithmetic progressions with union $\mathbf{Z}$ (so that $\sum_{i=1}^k \frac{1}{n_i} = 1$) contains two progressions with the same common difference. Henceforth we refer to any nontrivial finite disjoint family $\mathcal{F}$ of sets as a *tiling* of its union $\cup \mathcal{F}$ and speak of the sets in $\mathcal{F}$ as *tiles*. Generally speaking, one may ask for groups G that possess no tiling by cosets of distinct subgroups. Mirsky-Newman says $G = \mathbf{Z}$ constitutes one such group, but examples we give below show that the groups $\mathbf{Z}^d$ for $d > 2$ do not have this property (the case $d = 2$ remains open).

We now frame our positive result extending Mirsky-Newman to $\mathbf{Z}^d$, and obtained by restricting the permissible tiles to cosets of certain special subgroups.¹ We call any coset $T = \mathbf{v} + L$ of a subgroup L of $\mathbf{Z}^d$ ($d \geq 1$) a *Cartesian coset* if L has the form $a_1\mathbf{Z} \times \cdots \times a_d\mathbf{Z}$ for positive integers $a_1, \dots, a_d$. Thus a Cartesian coset in $\mathbf{Z}^d$ equals the Cartesian product of d arithmetic progressions. (Note that L must have full rank.)

THEOREM: $\mathbf{Z}^d$ admits no tiling using only Cartesian cosets no two of which are translates of one another.

Ahead of the proof, we gather a few facts of Fourier analysis. For $\mathbf{C}$, the complex numbers, every $\mathbf{C}$ -valued function on a finite abelian group equals a unique finite $\mathbf{C}$ -linear combination of characters (i.e. homomorphisms to $\mathbf{C}^*$). For L a full-rank subgroup of $\mathbf{Z}^d$, we may regard L -periodic functions on $\mathbf{Z}^d$ as functions on $\mathbf{Z}^d/L$ which thus expand as $\mathbf{C}$ -linear combinations of characters. These characters on $\mathbf{Z}^d/L$ pull back to $\mathbf{Z}^d$ as exponential functions $\mathbf{x} \mapsto \exp(2\pi i \mathbf{k} \cdot \mathbf{x})$ with $\mathbf{k}$ such that $\mathbf{k} \cdot L \subseteq \mathbf{Z}^d$. Adding an element of $\mathbf{Z}^d$ to $\mathbf{k}$ will not affect the exponential, so we standardize the components of $\mathbf{k}$ to lie in $[0, 1)$. Given L , we write $\widehat{L}$ for the associated finite set of standardized vectors $\mathbf{k}$. E.g., if $L = 2\mathbf{Z} \times 3\mathbf{Z}$ we have $\widehat{L} = \{(0, 0), (0, 1/3), (0, 2/3), (1/2, 0), (1/2, 1/3), (1/2, 2/3)\}$. Note that the cardinality of $\widehat{L}$ is precisely $[\mathbf{Z}^d : L]$.

Given an L -periodic function $f : \mathbf{x} \mapsto \sum_{\mathbf{k} \in \widehat{L}} c_{\mathbf{k}} \exp(2\pi i \mathbf{k} \cdot \mathbf{x})$, write $\widehat{f}(\mathbf{k}) = c_{\mathbf{k}}$. One can write $\widehat{f}(\mathbf{k})$ as $\frac{1}{[\mathbf{Z}^d : L]} \sum_{\mathbf{x}} f(\mathbf{x}) \exp(-2\pi i \mathbf{k} \cdot \mathbf{x})$ where $\mathbf{x}$ varies over a set of coset representatives. If f is L -periodic, then $\widehat{f}(\mathbf{k})$ vanishes for $\mathbf{k} \notin \widehat{L}$.

As usual, for each $S \subseteq \mathbf{Z}^d$ we have an indicator function χ_S , with $\chi_S(v) = 1$ if $v \in S$, and $= 0$ otherwise.

By symmetry, the sum of all characters on $\mathbf{Z}^d/L$ vanishes everywhere except the identity, where the value equals the cardinality of $\mathbf{Z}^d/L$; thus summing the corresponding exponentials and dividing by the index $[\mathbf{Z}^d : L]$ of L in $\mathbf{Z}^d$ yields χ_L . So $\widehat{\chi_L} = 1/[\mathbf{Z}^d : L]$ on $\widehat{L}$, 0 elsewhere.

$\chi_{\mathbf{v}+L}(\mathbf{x}) = \chi_L(\mathbf{x} - \mathbf{v})$, so $\widehat{\chi_{\mathbf{v}+L}}(\mathbf{k}) = (1/[\mathbf{Z}^d : L]) \exp(-2\pi i \mathbf{k} \cdot \mathbf{v})$ on $\widehat{L}$, 0 elsewhere.

PROOF: Suppose that $\mathbf{Z}^d = \cup T_j$ gives a tiling by Cartesian cosets $T_j =$

¹Where we strengthen the hypothesis, Sun's theorem generalizes Mirsky-Newman to $\mathbf{Z}^d$ by weakening the conclusion: every tiling of $\mathbf{Z}^d$ contains two tiles, cosets respectively of subgroups L and L' giving rise to isomorphic quotients of $\mathbf{Z}^d$. Whether or not we can require that L and L' differ merely by a rotation (in $O(d)$) we do not know.

$\mathbf{v}_j + L_j$. Tiling implies $\sum_j \chi_{T_j} = \chi_{\mathbf{Z}^d}$ and thus $\sum_j \widehat{\chi_{T_j}} = \widehat{\chi_{\mathbf{Z}^d}}$. For $L_j = a_1 \mathbf{Z} \times \cdots \times a_d \mathbf{Z}$, $\mathbf{k}_j := (1/a_1, \dots, 1/a_d) \in \widehat{L_j}$. Take L_m with $a_1 \cdots a_d$ maximal. Then $\widehat{\chi_{T_m}}(\mathbf{k}_m) \neq 0$ even though $\widehat{\chi_{\mathbf{Z}^d}}(\mathbf{k}_m) = 0$. So $\widehat{\chi_{T_j}}(\mathbf{k}_m) \neq 0$ for some $j \neq m$. Our choice of L_m implies $L_j = L_m$. $\square$

NOTE: We know no previous Fourier-theoretic proof of Mirsky-Newman. However, Henry Cohn has pointed out that in the case $d = 1$, the Fourier coefficients in our proof are precisely the residues at the poles of the terms in the partial fraction decomposition that occurs in the Mirsky-Newman power series proof. Indeed, the link between Fourier decomposition and partial fraction decomposition is discussed in chapter 7 of the book of Beck and Robins. This link extends to higher dimensions, and we intend to write a follow-up article that re-proves the main result of the present paper using d -variable generating functions. For a preview, see the Feldman-Propp-Robins preprint listed in the References.

The four cosets

$$(0, 1, 0) + 2\mathbf{Z} \times 2\mathbf{Z} \times \mathbf{Z},$$

$$(0, 0, 1) + \mathbf{Z} \times 2\mathbf{Z} \times 2\mathbf{Z},$$

$$(1, 0, 0) + 2\mathbf{Z} \times \mathbf{Z} \times 2\mathbf{Z}, \text{ and}$$

$$(2\mathbf{Z} \times 2\mathbf{Z} \times 2\mathbf{Z}) \cup ((1, 1, 1) + 2\mathbf{Z} \times 2\mathbf{Z} \times 2\mathbf{Z})$$

tile $\mathbf{Z}^3$, and no two of them are translates of one another; this shows that for $d = 3$, the Theorem becomes false if the Cartesian requirement is dropped. Furthermore, taking this example $\times \mathbf{Z}^{d-3}$ gives a d -dimensional counterexample for every $d > 3$.

ACKNOWLEDGMENTS: Conversations with Henry Cohn and suggestions from the referee were helpful in the preparation of this article for publication. The second author is supported by a grant from the National Science Foundation, and the third author is supported in part by Nanyang Technological University's SUG grant M5811053.

REFERENCES

- M. Beck and S. Robins, Computing the continuous discretely: integer-point enumeration in polyhedra, Springer, 2007.
P. Erdős, Egy kongruenciarendszerekről szóló problémáról (On a problem concerning covering systems). Mat. Lapok 3 (1952), 122–128 (Hungarian).
D. Feldman, J. Propp, and S. Robins, Tiling Lattices with Sublattices, II, arXiv: 1006.0472.

- D.J. Newman, *Analytic Number Theory*, Springer 2000.
- A. Soifer, *The Mathematical Coloring Book*, Springer 2008.
- Z. Sun, On the Herzog-Schönheim conjecture for uniform covers of groups,
J. Algebra 273 (2004), 153–175.